

OPEN CORACOID TRANSFER (LATARJET) REHABILITATION PROGRAM
(TO BE GIVEN TO PHYSICAL THERAPIST)

	RANGE OF MOTION	SLING	THERAPEUTIC EXERCISE
PHASE I: 0-6 WEEKS POST-OP	Limit ER to passive 45° to protect subscapularis repair FE progress as tolerated	0-2 weeks: Worn at all times (day and night) Off for gentle exercise only 2-6 weeks: Worn daytime only	0-3 weeks: Grip strengthening, pendulum exercises Elbow/wrist/hand ROM at home 3-6 weeks: Begin cuff, deltoid isometrics; limit ER to passive 45° No active IR nor extension until 6 weeks
PHASE II: 6-12 WEEKS POST-OP	Increase as tolerated to full Begin active assisted/active internal rotation and extension as tolerated after 6 weeks	None	6-8 weeks: Begin light resisted ER, forward flexion and abduction 8-12 weeks: Begin resisted internal rotation, extension and scapular retraction
PHASE III: 12-24 WEEKS POST-OP	Progress to full ROM without discomfort	None	Advance strengthening as tolerated Closed chain scapular rehab and functional rotator cuff strengthening; focus on anterior deltoid and teres Maximize subscapular stabilization